

Math 525, Differential Equations
Winter 2015
Assignment 3, due February 25 , 2015, 9 AM

Exercise 11: (2)

Find the spectrum and the corresponding eigenfunctions in $L^2([0, \pi])$ for the operators A and B with

$$A = -\frac{d^2}{dx^2}, \quad \mathcal{D}(A) = \{f \in L^2(0, \pi); Af \in L^2(0, \pi), f(0) = 0, f(\pi) = 0\},$$
$$B = -\frac{d^2}{dx^2}, \quad \mathcal{D}(B) = \{f \in L^2(0, \pi); Bf \in L^2(0, \pi), \frac{d}{dx}f(0) = 0, \frac{d}{dx}f(\pi) = 0\}.$$

Exercise 12: (2)

Let A be a symmetric linear operator on a Hilbert space H with $R(A) = H$ and with compact inverse A^{-1} . The natural domain of definition is

$$\mathcal{D}(A) = \{u \in H; Au \in H\}$$

Show that there exists an orthonormal basis $\{w_j\}$ of H such that

$$\mathcal{D}(A) = \left\{ u; u = \sum c_j w_j, \quad \sum |c_j|^2 \lambda_j^2 < \infty \right\}.$$

Exercise 13: (3)

For $A = -\Delta$ show that on a bounded domain the norm on $\mathcal{D}(A^{\frac{1}{2}})$ is equivalent to the norm on H^1 . (Hint: If A is symmetric, $(Au, v) = (u, Av)$, then also $A^{\frac{1}{2}}$).

Exercise 14: (3)

Show that if Ω is bounded, then $L^2(\Omega)$ is compactly embedded in $H^{-1}(\Omega)$.

Exercise 15: (5)

Let V be the subspace of $H^1(\Omega)$, consisting of functions with zero integral

$$V := \left\{ u \in H^1(\Omega) : \int_{\Omega} u(x) dx = 0 \right\}.$$

Arguing by contradiction, use Rellich-Kondrachov compactness to show that there exists a constant $C > 0$ such that we have a Poincaré inequality

$$\|u\|_2 \leq C \|\nabla u\|_2.$$

(Hint, you may assume that if $\nabla u = 0$ a.e., then $u = \text{const. a.e.}$).

Exercise 16: (The Nagumo Model) (5)

Fitzhugh and Nagumo derived a model for signal transduction in the axon:

$$u_t = u_{xx} + u(1-u)(u-\alpha),$$

where u represents the membrane potential, and $0 < \alpha < 1$. We study this model on $0 \leq x \leq l$ with homogeneous Neumann boundary conditions,

$$u_x(t, 0) = 0, \quad u_x(t, l) = 0.$$

- Find a system of two ODEs which describe the steady states. We denote this system of ODE's by (*) for now.
 - Study the steady states of (*) and their stability.
- (c)-(e) on next page.

(c) Show that

$$H(u, v) = \frac{1}{2}(u_x)^2 - \frac{1}{4}u^4 + \frac{1+\alpha}{3}u^3 - \frac{\alpha}{2}u^2$$

is a Hamiltonian function for the system (*).

- (d) Sketch the phase portrait of (*) in the (u, u_x) -plane. Distinguish the qualitatively different cases depending on $\alpha \in (0, 1)$.
- (e) Find the steady states which satisfy the Neumann boundary conditions, and plot the steady states u as a function of x .