

## UNIQUENESS AND ASYMPTOTICS

In this lecture we prove the uniqueness for the wave equations. We also prove some asymptotic decay results.

**1. Uniqueness via energy method.**

Consider the wave equation in a bounded domain  $\Omega \subset \mathbb{R}^n$ .

$$\square u \equiv u_{tt} - \Delta u = f \quad \Omega \times (0, T) \quad (1)$$

$$u = g \quad \Omega \times \{0\} \text{ and } \partial\Omega \times [0, T] \quad (2)$$

$$u_t = h \quad \Omega \times \{0\}. \quad (3)$$

It is clear that the uniqueness of this problem is equivalent to that the following equation

$$\square u \equiv u_{tt} - \Delta u = 0 \quad \Omega \times (0, T) \quad (4)$$

$$u = 0 \quad \Omega \times \{0\} \text{ and } \partial\Omega \times [0, T] \quad (5)$$

$$u_t = 0 \quad \Omega \times \{0\}. \quad (6)$$

having only 0 solution.

Now we prove this. Multiply the equation by  $u_t$  and integrate over  $\Omega \times (0, T)$ , we have

$$\begin{aligned} 0 &= \int_{\Omega \times (0, T)} (u_{tt} - \Delta u) u_t \, dx \, dt \\ &= \int_{\Omega \times (0, T)} \frac{d}{dt} \left( \frac{1}{2} u_t^2 \right) \, dx \, dt + \int_{\Omega \times (0, T)} -\Delta u u_t \, dx \, dt \\ &= \int_{\Omega \times (0, T)} \frac{d}{dt} \left( \frac{1}{2} u_t^2 \right) \, dx \, dt + \int_{\Omega \times (0, T)} \nabla u \cdot \nabla u_t \, dx \, dt \\ &= \int_{\Omega \times (0, T)} \frac{d}{dt} \left[ \frac{1}{2} (u_t^2 + |\nabla u|^2) \right] \, dx \, dt \\ &= \int_{\Omega} \left[ \frac{1}{2} (u_t^2 + |\nabla u|^2) \right] (x, T) \, dx - \int_{\Omega} \left[ \frac{1}{2} (u_t^2 + |\nabla u|^2) \right] (x, 0) \, dx \\ &= \int_{\Omega} \left[ \frac{1}{2} (u_t^2 + |\nabla u|^2) \right] (x, T) \, dx. \end{aligned} \quad (7)$$

This implies  $u$  is a constant at time  $T$ . But this constant must be 0 according to the boundary value.

**Remark 1.** If we know the solutions decays at infinity, we can use the same method when  $\Omega$  is unbounded and obtain the same result.

**2. Domain of dependence.**

We have seen from the formulas that the value of  $u(x, t)$  only depends on the initial values in the ball  $B_t(x)$ . In other words, if  $g = h = 0$  in  $B_r(x)$ , then  $u$  must vanish in the cone

$$|x| + t \leq r. \quad (8)$$

We prove this fact now.

Denote by  $C_r$  the above mentioned cone. and for  $T < r$  denote by  $U_T$  the following domain

$$U_T \equiv \{(x, t) \in C_r, 0 \leq t \leq T\}. \quad (9)$$

Then naturally the boundary of  $U_T$  consists of three parts

$$\partial U_T = S_T + S_0 + S_{\text{side}} \quad (10)$$

where

$$S_u = \{(x, t) \in C_r, t = u\}, \quad S_{\text{side}} = \partial C_r \cap \bar{U}_T. \quad (11)$$

Now we compute

$$\begin{aligned}
0 &= \int_{U_T} (u_{tt} - \Delta u) u_t \, dx \, dt \\
&= \int_{U_T} u_{tt} u_t - \Delta u u_t \, dx \, dt \\
&= \int_{U_T} \partial_t \left( \frac{1}{2} u_t^2 \right) - \nabla \cdot (\nabla u u_t) + \nabla u \cdot \nabla u_t \, dx \, dt \\
&= \int_{U_T} \partial_t \left( \frac{1}{2} u_t^2 + \frac{1}{2} |\nabla u|^2 \right) - \nabla \cdot (\nabla u u_t) \, dx \, dt \\
&= \int_{U_T} \nabla_{t,x} \cdot \begin{pmatrix} \frac{1}{2} u_t^2 + \frac{1}{2} |\nabla u|^2 \\ -\nabla u u_t \end{pmatrix} \, dx \, dt \\
&= \int_{\partial U_T} n_{t,x} \cdot \begin{pmatrix} \frac{1}{2} u_t^2 + \frac{1}{2} |\nabla u|^2 \\ -\nabla u u_t \end{pmatrix} \, dS \\
&= \int_{S_T} \frac{1}{2} u_t^2 + \frac{1}{2} |\nabla u|^2 - \int_{S_0} \frac{1}{2} u_t^2 + \frac{1}{2} |\nabla u|^2 \\
&\quad + \int_{S_{\text{side}}} \begin{pmatrix} n_t \\ n_x \end{pmatrix} \cdot \begin{pmatrix} \frac{1}{2} u_t^2 + \frac{1}{2} |\nabla u|^2 \\ -\nabla u u_t \end{pmatrix} \, dS \\
&= \int_{S_T} \frac{1}{2} u_t^2 + \frac{1}{2} |\nabla u|^2 - \int_{S_0} \frac{1}{2} u_t^2 + \frac{1}{2} |\nabla u|^2 \\
&\quad + \int_{S_{\text{side}}} \frac{n_t}{2} u_t^2 + \frac{n_t}{2} |\nabla u|^2 - n_x \cdot \nabla u u_t \, dS. \tag{12}
\end{aligned}$$

For the last term, we notice that the equation for  $S_{\text{side}}$  is  $|x| + t = r$  which means  $n_t = |n_x|$  and consequently

$$\frac{n_t}{2} u_t^2 + \frac{n_t}{2} |\nabla u|^2 - n_x \cdot \nabla u u_t \geq 0. \tag{13}$$

Thus we have shown that

$$\int_{S_T} \frac{1}{2} u_t^2 + \frac{1}{2} |\nabla u|^2 \leq \int_{S_0} \frac{1}{2} u_t^2 + \frac{1}{2} |\nabla u|^2 \tag{14}$$

for all  $T < r$  and the conclusion follows.

### 3. Decay of the solution.

We prove the following.

**Proposition 2.** *Let  $u$  solve*

$$u_{tt} - \Delta u = 0 \quad \text{in } \mathbb{R}^3 \times (0, \infty) \tag{15}$$

$$u = g \quad u_t = h \quad \text{on } \mathbb{R}^3 \times \{t=0\} \tag{16}$$

where  $g, h$  are smooth and have compact support. Then there is a constant  $C$  such that

$$|u(x, t)| \leq C/t \tag{17}$$

for all  $(x, t)$ .

**Proof.** Recall the Kirchhoff formula:

$$u(x, t) = \frac{1}{4\pi t^2} \int_{\partial B_t(x)} t h(w) + g(w) + \nabla g(w) \cdot (y - x) \, dS_w. \tag{18}$$

Since  $h, g, \nabla g$  vanishes outside their respective supports, we can write

$$u(x, t) = \frac{1}{4\pi t^2} \int_{\partial B_t(x) \cap A} t h(w) + g(w) + \nabla g(w) \cdot (y - x) \, dS_w \tag{19}$$

where  $A$  is the union of the three supports. Now the conclusion easily follows after we notice that the area of  $\partial B_t(x) \cap A$  is bounded by a constant independent of  $t$ .  $\square$

**Remark 3.** The above estimate behaves badly when  $t$  is small. But this is easily remedied by noticing that when  $t$  is small, the area of  $\partial B_t(x) \cap A$  scales as  $t^2$  and therefore  $u$  is uniformly bounded. Integrating this observation into the estimate gives

$$|u(x, t)| \leq C(1+t)^{-1}. \quad (20)$$

**Proposition 4.** *Let  $u$  solve*

$$u_{tt} - \Delta u = 0 \quad \text{in } \mathbb{R}^2 \times (0, \infty) \quad (21)$$

$$u = g \quad u_t = h \quad \text{on } \mathbb{R}^2 \times \{t=0\} \quad (22)$$

where  $g, h$  are smooth and have compact support. Then there is a constant  $C$  such that

$$|u(x, t)| \leq C(1+t)^{-1/2}(1+|t-|x||)^{-1/2}. \quad (23)$$

for all  $(x, t)$ .

**Proof.** Assume that the supports of  $g, h$  are contained in the ball  $B_R$ . Recall the Poisson's formula:

$$u(x, t) = \frac{1}{2\pi t^2} \int_{D_t(x)} \frac{t g(y) + t^2 h(y) + t \nabla g \cdot (y-x)}{(t^2 - |y-x|^2)^{1/2}} dy \quad (24)$$

By taking the supreme of  $g, h, \nabla g$  and noticing that  $|y-x| \leq t$  we have

$$|u(x, t)| \leq C \int_{D_t(x)} \frac{dy}{(t-|y-x|)^{1/2}(t+|y-x|)^{1/2}} \leq C t^{-1/2} \int_{D_t(x)} \frac{dy}{(t-|y-x|)^{1/2}}. \quad (25)$$

Now let  $z = y - x$  we have

$$|u(x, t)| \leq C t^{-1/2} \int_{D_t} \frac{dz}{(t-|z|)^{1/2}}. \quad (26)$$

Here note that the integral is in fact over  $D_t \cap \{|z+x| \leq R\}$ . We have

- $|x| > t + R$ :  $u(x, t) \equiv 0$ .
- $t - 2R < |x| < t + R$ : We use polar coordinates, note that the angle is of order  $R/t$  (we only consider the case  $t \gg R$  here), thus we have

$$\begin{aligned} \int_{D_t \cap \{|z+x| \leq R\}} \frac{dz}{(t-|z|)^{1/2}} &\lesssim \frac{R}{t} \int_{|x|-R}^{\min(t, |x|+R)} \frac{r dr}{(t-r)^{1/2}} \\ &\leq R \int_{|x|-R}^{\min(t, |x|+R)} (t-r)^{-1/2} dr \\ &\leq C(t-(|x|-R))^{1/2} \leq C. \end{aligned} \quad (27)$$

- $|x| < t - 2R$ : We have  $(t-|z|) \geq c(1+t-|x|)$ , and therefore

$$\int_{D_t \cap \{|z+x| \leq R\}} \frac{dz}{(t-|z|)^{1/2}} \leq C(1+t-|x|)^{-1/2} |B_R|. \quad (28)$$

Combining the above, we see that when  $t$  is large (for example  $t > 3R$ ), we have

$$|u(x, t)| \leq C t^{-1/2}(1+|t-|x||)^{-1/2}. \quad (29)$$

When  $t \leq 3R$ , we have

$$\begin{aligned} |u(x, t)| &\leq C \int_{D_t} \frac{dz}{(t^2 - |z|^2)^{1/2}} \\ &= C \int_0^t \frac{r dr}{(t^2 - r^2)^{1/2}} \\ &= C t \leq 3CR. \end{aligned} \quad (30)$$

Thus  $u$  is bounded by a constant when  $t \leq 3R$  and by  $Ct^{-1/2}(1 + |t - |x||)^{-1/2}$  when  $t > 3R$ , as a consequence, we can write

$$|u(x, t)| \leq C(1+t)^{-1/2}(1 + |t - |x||)^{-1/2}. \quad (31)$$

as desired.  $\square$

**Remark 5.** In general, we have

–  $n$  odd:

$$|u(t, x)| \leq C(1+t)^{-\frac{n-1}{2}}; \quad (32)$$

–  $n$  even:

$$|u(t, x)| \leq C(1+t)^{-\frac{n-1}{2}}(1 + |t - |x||)^{-\frac{n-1}{2}}. \quad (33)$$

**Remark 6.** Such algebraic decays are also characteristic in other dispersive equations, for example the Schrödinger equation.

**Remark 7.** It is clear that no decay can be expected for the solutions to the 1D wave equation:

$$u(x, t) = \frac{1}{2}[g(x+t) + g(x-t)] + \frac{1}{2} \int_{x-t}^{x+t} h(y) dy. \quad (34)$$

### Exercises.

**Exercise 1.** Consider the 1D wave equation with variable coefficients  $a(x, t) \in C^\infty(\mathbb{R}^n \times [0, \infty))$  with  $a(x, t) > a_0 > 0$  for some constant  $a_0$

$$u_{tt} - a(x)^2 u_{xx} = 0. \quad (35)$$

Find out and prove its domain of dependence.