The extension of the finite-dimensional version of Krivine's theorem to quasi-normed spaces.

A.E. Litvak*

Recently, a number of results of the Local Theory have been extended to the quasi-normed spaces. There are several works ([Kal1], [Kal2], [D], [GL], [KT], [GK], [BBP1], [BBP2], [M2]) where such results as Dvoretzky-Rogers lemma ([DvR]), Dvoretzky theorem ([Dv1], [Dv2]), Milman's subspace-quotient theorem ([M1]), Krivine's theorem ([Kr]), Pisier's abstract version of Grotendick's theorem ([P1], [P2]), Gluskin's theorem on Minkowski compactum ([G]), Milman's reverse Brunn-Minkowski inequality ([M3]), and Milman's isomorphic regularization theorem ([M4]) are extended to quasi-normed spaces after they were established for normed spaces. It is somewhat surprising since the first proofs of these facts substantially used convexity and duality.

In [AM2] D. Amir and V.D. Milman proved the local version of Krivine's theorem (see also [Gow], [MS]). They studied quantitative estimates appearing in this theorem. We extend their result to the q- and quasi-normed spaces.

Recall that the quasi-norm on a real vector space X is a map $\|\cdot\| : X \longrightarrow \mathbb{R}^+$ such that
1) $\|x\| > 0 \quad \forall x \neq 0$,
2) $\|tx\| = |t| \, \|x\| \quad \forall t \in \mathrm{R}, \ x \in X$,
3) $\exists C \geq 1$ such that $\forall x, y \in X \ \ \|x+y\| \leq C(\|x\| + \|y\|)$.
If 3) is substituted by
3a) $\forall x, y \in X \ \|x+y\|^q \leq \|x\|^q + \|y\|^q \quad$ for some fixed $q \in (0,1]$
then $\|\cdot\|$ is called a q-norm on X. Note that 1-norm is the usual norm. It is obvious that every q-norm is a quasi-norm with $C = 2^{1/q-1}$. However, not every quasi-norm is q-norm for some q. Moreover, it is even not necessary

*This research was supported by Grant No.92-00285 from United States-Israel Binational Science Foundation (BSF).

continuous. It can be shown by the following simple example. Let f be a positive function on the Euclidean sphere S^{n-1} defined by

$$f(x) = \begin{cases} |x| & \text{for } x \in A, \\ 2|x| & \text{otherwise.} \end{cases}$$

Here A is a subset of S^{n-1} such that both A and $S^{n-1} \setminus A$ are dense in S^{n-1}. Denote $\|x\| = |x| f(x/|x|)$. Because f is not continuous it is clear that $\|\cdot\|$ is not q-norm for any q though it is the quasi-norm.

The next lemma is Aoki-Rolewicz Theorem ([KPR], [R], see also [K], p.47).

Lemma 1 *Let $\|\cdot\|$ be a quasi-norm with the constant C in the quasi-triangle inequality. Then there exists a q-norm $\|\cdot\|$ for which*

$$\|x\|_q \le \|x\| \le 2C\|x\|_q$$

with q satisfying $2^{1/q-1} = C$. This q-norm can be defined as follows

$$\|x\|_q = \inf\left\{ \left(\sum_{i=1}^n \|x_i\|^q\right)^{1/q} : n > 0,\ x = \sum_{i=1}^n x_i \right\}.$$

We refer to [KPR] for further properties of the quasi- and q-norms.

Next, we prove the following theorem.

Theorem 1 *Let $\{e_i\}_1^n$ be a unit vector basis in $\mathbb{R}^n$, $\|\cdot\|_p$ be a l_p-norm on $\mathbb{R}^n$, i.e. $\|\sum_{i=1}^n a_i e_i\|_p = (\sum_i |a_i|^p)^{1/p}$, for $0 < p < \infty$. Let $\|\cdot\|$ be a q-norm on $\mathbb{R}^n$ such that*

$$C_1^{-1}\|x\|_p \ \le\ \|x\| \le C_2\|x\|_p \qquad (*)$$

for every $x \in \mathbb{R}^n$. Then for every $\varepsilon > 0$ and $C = C_1 C_2$ there exists a block sequence $u_1, u_2, ..., u_m$ of $e_1, e_2, ..., e_n$ which satisfies

$$(1-\varepsilon)\left(\sum_{i=1}^m |a_i|^p\right)^{1/p} \ \le\ \left\|\sum_{i=1}^m a_i u_i\right\| \ \le\ (1+\varepsilon)\left(\sum_{i=1}^m |a_i|^p\right)^{1/p} \qquad (**)$$

for all $a_1, a_2, ..., a_m$ and $m \ge C(\varepsilon, p, q)\,(n/\log n)^\nu$, where

$$\nu = \frac{\alpha\varepsilon_0}{\varepsilon_0 + p + \alpha\varepsilon_0}\,,\ \text{for}\ \ p < 1\ \ \text{and}\ \ \nu = \frac{\varepsilon_0}{2\varepsilon_0 + 1}\,,\ \text{for}\ \ p \ge 1\,;$$

$$\alpha = \min\{p, q\}\,,\ \varepsilon_0 = \left(\frac{q\varepsilon/2}{1 + C^q 12^{q/p}}\right)^{p/q}.$$

Remark 1. If $p \geq 1$ in this theorem, then we have the well-known finite-dimensional version of Krivine's theorem with some modifications concerning change of the usual norm to the q-norm. In this case for small enough q we get $\varepsilon_0 \approx \left(\frac{q\varepsilon}{4}\right)^{p/q}$ and $\nu \approx \varepsilon_0$.

The case $p < 1$ is more interesting. We get an extension of the finite-dimensional version of Krivine's theorem. To provide an intuition for the behavior of the constant in the theorem we point out that for small enough p and q with $p = q$ we can take $\varepsilon_0 \approx \frac{q\varepsilon}{30}$ and $\nu \approx \varepsilon_0$.

Remark 2. By Lemma 1 in the case of quasi-norm with the constant C_0 the inequality $(**)$ is substituted with

$$(1-\varepsilon)\left(\sum_{i=1}^{m}|a_i|^p\right)^{1/p} \leq \left\|\sum_{i=1}^{m}a_iu_i\right\| \leq 2(1+\varepsilon)C_0\left(\sum_{i=1}^{m}|a_i|^p\right)^{1/p}.$$

Due to the example above, we can not remove the constant C_0 in this inequality.

The proof of the theorem consists of two lemmas.

Lemma 2 *For every $\eta > 0$ there exists a constant $C(\eta) > 0$ such that if $\|\cdot\|$ is a q-norm on $\mathbb{R}^n$ satisfying $(*)$ then there exists a block sequence $y_1, y_2, ..., y_k$ of $e_1, e_2, ..., e_n$ which is $(1+\eta)$-symmetric and $k \geq C(\eta, q, p)\frac{n}{\log n}$.*

Lemma 3 *If $y_1, y_2, ..., y_k$ is a 1-symmetric sequence in a normed space satisfying*

$$C_1^{-1}\|a\|_p \leq \left\|\sum_{i=1}^{k}a_iy_i\right\| \leq C_2\|a\|_p$$

for all $a = (a_1, a_2, ..., a_k) \in \mathbb{R}^k$ then for every $\varepsilon > 0$ there exists a block sequence $u_1, u_2, ..., u_m$ of $y_1, y_2, ..., y_k$ such that

$$(1-\varepsilon)\|a\|_p \leq \left\|\sum_{i=1}^{m}a_iu_i\right\| \leq (1+\varepsilon)\|a\|_p$$

for all $a = (a_1, a_2, ..., a_m) \in \mathbb{R}^m$, where $m \geq C(p,q)\varepsilon^{p/q}k^{\nu}$, $\nu = \frac{\alpha\varepsilon_0}{\varepsilon_0+p+\alpha\varepsilon_0}$, for $p < 1$ and $\nu = \frac{\varepsilon_0}{2\varepsilon_0+1}$, for $p \geq 1$, $\alpha = \min\{p, q\}$, $\varepsilon_0 = \left(\frac{q\varepsilon}{1+C^q12^{q/p}}\right)^{p/q}$.

At first, D. Amir and V.D. Milman ([AM2], see also [MS]) proved Lemma 2 for $q = 1$, $p \geq 1$ with the estimate $k \geq C(\eta, q, p)\, n^{1/3}$. Their proof can be modified to obtain result for $0 < p < \infty$, $q \leq 1$. Afterwards, W.T. Gowers ([Gow]) showed that the estimate of k can be improved to $k \geq C(\eta, q, p)n/\ln n$. In fact, he gave two different, though similar, proofs for cases $p = 1$ and $p > 1$. The proof given for case $p = 1$ strongly used the convexity of the norm and the fact that p is equal to 1. However, the method used for $p > 1$ actually works for every $0 < p < \infty$ and even for q-norms. Let us recall the idea of W.T. Gowers. First we will introduce some definition.

Let Ω be the group $\{-1, 1\}^n \times S_n$, where S_n is the permutation group. Let Ψ be the group $\{-1, 1\}^k \times S_k$. For

$$b = \sum_{i=1}^{n} b_i e_i \in I\!R^n, \quad a = \sum_{i=1}^{k} a_i e_i \in I\!R^k, \quad (\varepsilon, \pi) \in \Omega, \quad (\eta, \sigma) \in \Psi$$

denote

$$b_{\varepsilon\pi} = \sum_{i=1}^{n} \varepsilon_i b_i e_{\pi(i)}\,, \qquad a_{\eta\sigma} = \sum_{i=1}^{k} \eta_i a_i e_{\sigma(i)}\,.$$

Let $h \cdot k = n$. For $i \leq k$, $j \leq h$ put

$$e_{ij} = e_{(i-1)h+j}\,, \quad \varepsilon_{ij} = \varepsilon_{(i-1)h+j}\,, \quad \pi_{ij} = \pi((i-1)h + j).$$

Define an action of Ψ on Ω by

$$\Psi_{\eta\sigma}((\varepsilon, \pi)) = (\varepsilon^1, \pi^1), \text{ where } \varepsilon^1_{ij} = \eta_i \varepsilon_{\sigma(i)j}, \quad \pi^1_{ij} = \pi_{\sigma(i)j}.$$

For any $(\varepsilon, \pi) \in \Omega$ define the operator

$$\Phi_{\varepsilon\pi} : I\!R^k \longrightarrow I\!R^n \qquad \text{by} \qquad \Phi_{\varepsilon\pi}\left(\sum_{i=1}^{k} a_i e_i\right) = \sum_{i=1}^{k}\sum_{j=1}^{h} \varepsilon_{ij} a_i e_{\pi_{ij}}.$$

For every $a \in I\!R^k$ by M_a denote the median of $\Phi_{\varepsilon\pi}(a)$ taken over Ω. Finally, let $A = \{a \in l_p^k : \|a\|_p \leq 1, a_1 \geq a_2 \geq \ldots \geq a_k \geq 0\}$.

The following claim, which W.T. Gowers proved for case $p > 1$ and $q = 1$, is the main step in the proof of Lemma 2.

Claim 1 *Let $\|\cdot\|$ be a q-norm on $I\!R^n$ satisfying $\|x\|_p \leq \|x\| \leq B\|x\|_p$. There is a constant $C_0 = C(p, q, \delta, B)$ such that given $\lambda > 0$ for every $a \in A$*

$$\mathbf{Prob}_\Omega\left\{\exists(\eta, \sigma) \ : \ |\, \|\Phi_{\varepsilon\pi}(a_{\eta\sigma})\|^q - M_a^q\,|^{1/q} > \frac{1}{2^{1/q}}\delta\|a\|_p h^{1/p}\right\} < 1/N$$

with $k = C_0 \frac{n}{\lambda \log n}$ and $N = k^\lambda$.

The proof of this claim can be equally well applied for all $0 < p < \infty$ and $0 < q \leq 1$. The only change that we have to do is to replace the triangle inequality

$$| \, \|x\| - \|y\| \, | \leq \|x - y\| \quad \text{by} \quad | \, \|x\|^q - \|y\|^q \, |^{1/q} \leq \|x - y\| \, .$$

The following two claims are technical and can be proved using ideas of [Gow] with small changes, connected with replacing $p \geq 1$ by $p < 1$ and the norm by q-norm.

Claim 2 *Let $0 < p < \infty$ and $\delta > 0$. There exist a constant λ, depending on p and δ only, such that for every integer k the set A contains a δ-net K of cardinality k^λ.*

Claim 3 *Let $|| \cdot ||$ be a q-norm on $\mathbb{R}^n$ satisfying $\|x\|_p \leq \|x\| \leq B\|x\|_p$. If there is $(\varepsilon, \pi) \in \Omega$ such that for every a in some δ-net K of A*

$$| \, \|\Phi_{\varepsilon\pi}(a_{\eta\sigma})\|^q - \|\Phi_{\varepsilon\pi}(a_{\eta_1\sigma_1})\|^q \, |^{1/q} \leq \delta \|a\|_p h^{1/p}$$

for every $(\eta, \sigma), (\eta_1, \sigma_1) \in \Psi$ then the block basis

$$\{\Phi_{\varepsilon\pi}(e_i)\}_{i=1}^k$$

of $(\mathbb{R}^k, || \cdot ||)$ is $(1 + 6\,(B\delta)^q)^{1/q}$-symmetric.

These three claims imply Lemma 2 in the standard way (see [Gow] for the details).

Proof of Lemma 3:
Our method of proof is close to the method used in [AM1], but our notation follows that of [MS] (ch. 10).

First, we will give the Krivine's construction of block basis. Let a and N be some integers which will be specified later. Let us introduce some set of numbers $\{\lambda_j\}_J$. We will say that set

$$\{B_{j,i}\}_{j \in J, i \in I}$$

(if *card* $I = 1$ then we have only one index j) is $\{\lambda_j\}_J$-set if
1) $B_{j,i} \subset \{1, ..., n\}$ for every $j \in J, i \in I$,

2) $B_{j,i}$ are mutually disjoint,
3) $card\ B_{j,i} = \lambda_j$ for every $j \in J,\ i \in I$.

Let us fix some $\{[\rho^j]\}$-set

$$\{A_{j,s}\}_{0 \le j \le N-1, 1 \le s \le m}$$

for $\rho = 1 + 1/a$.

For $0 \le j \le N-1$, $1 \le s \le m$ denote

$$Y_{j,s} = \sum_{i \in A_{j,s}} y_i$$

and define

$$z_s = \sum_{j=0}^{N-1} \rho^{(N-j)/p} Y_{j,s}.$$

Clearly, $\|z_1\| = \|z_2\| = ... = \|z_m\|$. The integer m will be defined from

$$k \approx m \sum_{j=0}^{N-1} [\rho^{(N-j)/p}] \approx m\rho^N(\rho-1)^{-1} = ma\left(\frac{a+1}{a}\right)^N.$$

Finally, we define the block sequence $\{u_s\}_{s=1}^m$ by

$$u_s = z_s/\|z_s\|.$$

Now, as in [MS], we will establish the necessary estimates.

Fix $N, M \in \{T+1, T+2, ..., m\}$ and $t_s \in \{0, ..., T\}$ for $s \in \{1, ..., m\}$ such that

$$\sum_{s=1}^{M} \rho^{-t_s} = 1 + \eta,\ |\eta| = 1.$$

Then

$$\sum_{s=1}^{M} \rho^{-t_s/p} z_s = \sum_{s=1}^{M} \sum_{j=0}^{N-1} \rho^{(N-j-t_s)/p} Y_{j,s} =$$

$$= \sum_{i=0}^{N-1+T} \rho^{(N-i)/p} \sum_{s \le M,\, j \le N-1,\, j+t_s=i} \ \sum_{l \in A_{j,s}} y_l = \sum_{i=0}^{N-1+T} \rho^{(N-i)/p} \sum_{l \in B_i} y_l$$

for some $\{a_i\}$-set $\{B_i\}_{i=0}^{N-1+T}$, where

$$a_i = \sum_{s \le M,\, j \le N-1,\, j+t_s=i} [\rho^{i-t_s}],\ 0 \le i \le N-1+T\,.$$

Therefore, we can choose a vector z which has the same structure as z_s (i.e. $z = \sum_{j=0}^{N-1} \rho^{(N-j)/p} \sum_{i\in A_j} y_i$ for some $\{[\rho^j]\}$-set $\{A_j\}_{0\le j\le N-1}$) such that the difference Δ is

$$\Delta = \sum_{s=1}^{M} \rho^{-t_s/p} z_s - z = \sum_{s=1}^{N-1} \rho^{(N-i)/p} \sum_{l\in C_i} y_l + \sum_{s=N}^{N-1+T} \rho^{(N-i)/p} \sum_{l\in C_i} y_l$$

for some $\{b_j\}$-set $\{C_j\}_{i=0}^{N-1+T}$, where

$$b_j = \begin{cases} |[\rho^j - a_j]| & \text{for } 0 \le j \le N-1, \\ a_j & \text{for } N \le j \le N-1+T. \end{cases}$$

Using technique of [MS] (pp. 66-67) we obtain

$$\|\Delta\| \le C_2 \rho^{N/p} (4T + N|\eta| + NM\rho^{-T})^{1/p} \quad \text{and} \quad \|z\| \ge (1/C_1)\rho^{N/p}(N/2)^{1/p}.$$

Hence

$$\left| \left\| \sum_{s=1}^{M} \rho^{-t_s/p} u_s \right\|^q - 1 \right| \le \left\| \sum_{s=1}^{M} \rho^{-t_s/p} u_s - \frac{z}{\|z\|} \right\|^q =$$

$$= \left(\frac{\|\Delta\|}{\|z\|} \right)^q \le (C_1 C_2)^q \Big(\frac{8T}{N} + 2|\eta| + 2M\rho^{-T} \Big)^{q/p}.$$

Thus

$$\left| \left\| \sum_{s=1}^{M} \rho^{-t_s/p} u_s \right\|^q - 1 \right| \le C^q (12\varepsilon_0)^{q/p},$$

provided $T \le N\varepsilon_0$, $|\eta| \le \varepsilon_0$, $M\rho^{-T} \le m\rho^{-T} \le \varepsilon_0$, for some ε_0.
Assume $T = [N\varepsilon_0]$.

CASE 1. $p < 1$.

Let $\sum_{s=1}^{m} |\alpha_s|^p = 1$ and $a_s = |\alpha_s|$. Let $\alpha = \min\{p, q\}$ and $\delta = \varepsilon_0^{1/p}/m^{1/\alpha}$. Take $\beta_s = \rho^{-t_s/p}$ or $\beta_s = 0$, $t_s \in \{0, 1, ..., T\}$ such that $|a_s - \beta_s| \le \delta$ for every s. It is possible if $\rho^{-T/p} \le \delta$ and $1 - \rho^{-1/p} \le \delta$. Since $p \le 1$ it is enough to take a such that it satisfies following the inequalities

$$\Big(\frac{a}{a+1} \Big)^{[N\varepsilon_0]} \le \delta^p = \frac{\varepsilon_0}{m^{p/\alpha}} \quad \text{and} \quad \delta \ge \frac{1}{p(a+1)}.$$

Take $a = \left[\frac{1}{\delta p}\right] = \left[\frac{m^{1/\alpha}}{p\varepsilon_0^{1/p}}\right]$. Thus $\delta \ge \frac{1}{p(a+1)}$,

$$\Big|\sum \rho^{-t_s} - 1\Big| = \Big|\sum \beta_s^p - 1\Big| \le \Big|\sum (a_s + \delta)^p - 1\Big| \le$$

$$\leq \left|\sum (a_s^p + \delta^p) - 1\right| = \delta^p m \leq \varepsilon_0\,,$$

and

$$\left| \left\|\sum_{s=1}^m \beta_s u_s\right\|^q - \left\|\sum_{s=1}^m \alpha_s u_s\right\|^q \right| \leq \left\|\sum_{s=1}^m |\beta_s - a_s| u_s\right\|^q \leq$$

$$\leq \delta^q \left\|\sum_{s=1}^m u_s\right\|^q \leq \delta^q m \leq \varepsilon_0^{q/p}\,.$$

Hence

$$\left| \left\|\sum_{s=1}^m \alpha_s u_s\right\|^q - 1 \right| \leq \varepsilon_0^{q/p}(1 + C^q 12^{q/p})\,,$$

if $m^{p/\alpha} \leq \varepsilon_0(\frac{1+a}{a})^{[N\varepsilon_0]}$ and $ma(\frac{1+a}{a})^N \leq k$, when $a = \left[\frac{m^{1/\alpha}}{p\varepsilon_0^{1/p}}\right]$. Choose N such that $(\frac{a}{1+a})^{N\varepsilon_0}$ is of the order $\varepsilon_0/m^{p/\alpha}$. Then

$$m\,\frac{m^{1/\alpha}}{p\varepsilon_0^{1/p}}\,\left(\frac{m^{p/\alpha}}{\varepsilon_0}\right)^{1/\varepsilon_0} = \frac{m^{1+1/\alpha+p/(\alpha\varepsilon_0)}}{\varepsilon_0^{1/p} p \varepsilon_0^{1/\varepsilon_0}} \sim k\,.$$

Thus, since $1/\alpha \geq \max\{1/p, 1/q\}$,

$$m \sim \varepsilon_0\,(pk)^{\frac{\alpha\varepsilon_0}{\varepsilon_0+p+\alpha\varepsilon_0}} \sim \varepsilon_0 k^{\frac{\alpha\varepsilon_0}{\varepsilon_0+p+\alpha\varepsilon_0}}$$

and for $\varepsilon_1 = \varepsilon_0^{q/p}\left(1 + c^q 12^{q/p}\right)$

$$(1-\varepsilon_1)^{1/q}\,\|(\alpha_s)\|_p \leq \left\|\sum \alpha_s u_s\right\| \leq (1+\varepsilon_1)^{1/q}\,\|(\alpha_s)\|_p$$

holds. For ε_1 small enough ($\varepsilon_1 < 2^q - 1$) we obtain $1 - \varepsilon_1/q \leq (1-\varepsilon_1)^{1/q}$ and $1 + 2\varepsilon_1/q \geq (1+\varepsilon_1)^{1/q}$. Take $\varepsilon = 2\varepsilon_1/q$, then

$$\varepsilon_0 = \left(\frac{q\varepsilon/2}{1 + C^q 12^{q/p}}\right)^{p/q}$$

and

$$m \geq C(p,q)\varepsilon^{p/q} k^{\frac{\alpha\varepsilon_0}{\varepsilon_0+p+\alpha\varepsilon_0}}\,.$$

CASE 2. $p \geq 1$**.**

We use the same idea. Let $\sum_{s=1}^m |\alpha_s|^p = 1$ and $a_s = |\alpha_s|$. Let $\delta = \varepsilon_0/(C^p m)$. Take $\beta_s = \rho^{-t_s/p}$ or $\beta_s = 0$, $t_s \in \{0, 1, ..., T\}$ such that $|a_s^p - \beta_s^p| \leq$

δ for every s. It is possible if $\rho^{-T} \le \delta$ and $1-\rho^{-1} \le \delta$. These two conditions are met if

$$\Big(\frac{a}{a+1}\Big)^{[N\varepsilon_0]} \le \delta = \frac{\varepsilon_0}{C^p m} \quad \text{and} \quad \delta \ge \frac{1}{a+1}\,.$$

Take $a = \left[\frac{1}{\delta}\right] = \left[\frac{C^p m}{\varepsilon_0}\right]$. Thus

$$\Big|\sum \rho^{-t_s} - 1\Big| = \Big|\sum \beta_s^p - 1\Big| \le \Big|\sum (a_s^p + \delta) - 1\Big| = \delta m \le \varepsilon_0.$$

Since

$$\Big\|\sum_{s=1}^m u_s\Big\| \le C_1 C_2 \frac{\|\sum_{s=1}^m u_s\|_p}{\|z\|_p} \le C_1 C_2 \Big(\frac{m \sum \rho^{N-j}[\rho^j]}{\|z\|_p^p}\Big)^{1/p} = C m^{1/p}$$

and

$$|\beta_s - a_s| \le |\beta_s^p - a_s^p|^{1/p} \le \delta^{1/p}\,,$$

we obtain

$$\Big|\ \Big\|\sum_{s=1}^m \beta_s u_s\Big\|^q - \Big\|\sum_{s=1}^m \alpha_s u_s\Big\|^q\Big| \le \Big\|\sum_{s=1}^m |\beta_s - a_s| u_s\Big\|^q \le$$

$$\le \delta^{q/p} \|\sum_{s=1}^m u_s\|^q \le \delta^{q/p} C^q m^{q/p} \le \varepsilon_0^{q/p}\,.$$

Hence

$$\Big|\ \Big\|\sum_{s=1}^m \alpha_s u_s\Big\|^q - 1\Big| \le \varepsilon_0^{q/p}(1 + C^q 12^{q/p})\,,$$

if $m \le \frac{\varepsilon_0}{C^p}(\frac{1+a}{a})^{[N\varepsilon_0]}$ and $ma(\frac{1+a}{a})^N \le k$, when $a = \left[\frac{C^p m}{\varepsilon_0}\right]$. Choose N such that $(\frac{a}{1+a})^{N\varepsilon_0}$ is of the order $\varepsilon_0/(C^p m)$. Then

$$m\ \frac{C^p m}{\varepsilon_0}\ \Big(\frac{C^p m}{\varepsilon_0}\Big)^{1/\varepsilon_0} = \Big(\frac{C^p}{\varepsilon_0}\Big)^{1+1/\varepsilon_0} m^{2+1/\varepsilon_0} \sim k\,.$$

Thus

$$m \ge \frac{\varepsilon_0}{C^p} k^{\frac{\varepsilon_0}{2\varepsilon_0+1}}$$

and for $\varepsilon_1 = \varepsilon_0^{q/p}\left(1 + C^q 12^{q/p}\right)$

$$(1-\varepsilon_1)^{1/q} \|(\alpha_s)\|_p \le \Big\|\sum \alpha_s u_s\Big\| \le (1+\varepsilon_1)^{1/q} \|(\alpha_s)\|_p$$

holds. For ε_1 small enough ($\varepsilon_1 < 2^q - 1$) we obtain $1 - \varepsilon_1/q \le (1-\varepsilon_1)^{1/q}$ and $1 + 2\varepsilon_1/q \ge (1+\varepsilon_1)^{1/q}$. Take $\varepsilon = 2\varepsilon_1/q$, then

$$\varepsilon_0 = \left(\frac{q\varepsilon/2}{1 + C^q 12^{q/p}}\right)^{p/q}$$

and

$$m \ge C(p,q)\varepsilon^{p/q} k^{\frac{\varepsilon_0}{2\varepsilon_0+1}} .$$

□

Acknowledgment. I want to thank Prof. V. Milman for his guidance and encouragement, and to thank the referee for his helpful remarks.

References

[**AM1**] D. Amir, V.D. Milman, *Unconditional and symmetric sets in n-dimensional normed spaces*, Isr. J. Math., 37, 3-20, 1980.

[**AM2**] D. Amir, V.D. Milman, *A quantitative finite-dimensional Krivine theorem*, Isr. J. Math., 50, 1-12, 1985.

[**BBP1**] J. Bastero, J. Bernués, A. Peña, *An extension of Milman's reverse Brunn-Minkowski inequality*, GAFA, 5, no. 3, 572-581, 1995.

[**BBP2**] J. Bastero, J. Bernués and A. Peña, *The theorem of Caratheodory and Gluskin for* $0 < p < 1$, Proc. AMS, 123, 1, 141- 144, 1995.

[**D**] S.J. Dilworth, *The dimension of Euclidean subspaces of quasi-normed spaces*, Math. Proc. Camb. Phil. Soc., 97, 311-320, 1985.

[**Dv1**] A. Dvoretzky, *A theorem on convex bodies and applications to Banach spaces*, Proc.Nat.Acad.Sci. U.S.A. 45, 223-226, 1959.

[**Dv2**] A. Dvoretzky, *Some results on convex bodies and Banach spaces*, Proc. Symp. Linear spaces, Jerusalem, 123-160, 1961.

[**DvR**] A. Dvoretzky and C. Rogers, *Absolute and unconditional convergence in normed linear spaces*, Proc. Nat. Acad. Sci. U.S.A. 36, 192-197, 1950.

[**G**] E. Gluskin, *The diameter of the Minkowski compactum is approximately equal to n*, Functional Anal. Appl. 15, 72-73, 1981.

[**GK**] Y. Gordon, N.J. Kalton, *Local structure theory for quasi-normed spaces*, Bull. Sci. Math., 118, 441-453, 1994.

[**GL**] Y. Gordon, D.R. Lewis, *Dvoretzky's theorem for quasi-normed spaces*, Illinois J. Math., 35, 250-259, 1991.

[**Gow**] W.T. Gowers, *Symmetric block bases in finite-dimensional normed spaces*, Isr. J. Math., 68, 193-219, 1989.

[**Kal1**] N.J. Kalton, *The convexity type of a quasi-Banach space* (unpublished note), 1977.

[Kal2] N.J. Kalton, *Convexity, type and the three space problem*, Studia Math., Vol. 69, 247-287, 1981.

[KPR] N.J. Kalton, N.T. Peck, J.W. Roberts, *An F-space sampler*, London Mathematical Society Lecture Note Series, 89, Cambridge University Press, Cambridge-New York, 1984.

[KT] N. Kalton, Sik-Chung Tam, *Factorization theorems for quasi-normed spaces*, Houston J. Math., 19, 301-317, 1993.

[K] H. König, *Eigenvalue Distribution of Compact Operators*, Birkhäuser, 1986.

[Kr] J.L. Krivine, *Sous-espaces de dimension finie des espaces de Banach réticulés*, Ann. Math. 104, 1-29, 1976.

[M1] V.D. Milman, *Almost Euclidean quotient spaces of subspaces of a finite dimensional normed space*, Proc. Amer. Math. Soc. 94, no. 3, 445-449, 1985.

[M2] V. Milman, *Isomorphic Euclidean regularization of quasi-norms in $\mathbb{R}^n$*, C. R. Acad. Sci. Paris, 321, 879-884, 1996.

[M3] V.D. Milman, *Inégalité de Brunn-Minkowsky inverse et applications á la théorie locale des espaces normés*, C.R. Acad. Sci. Paris 302, Sér 1, 25-28, 1986.

[M4] V. Milman, *Some applications of duality relations*, GAFA Seminar 1989-90, Springer Lecture Notes in Math. 1469, 13-40, 1991.

[MS] V. Milman, G. Schechtman, *Asymptotic theory of finite dimensional normed spaces*, Springer Lecture notes 1200, 1986.

[P1] G. Pisier, *Un théorème sur les opérateurs entre espaces de Banach qui se factorisent par un espace de Hilbert*, Ann. École Norm. Sup. Vol. 13, 23-43, 1980.

[P2] G. Pisier, *Factorization of linear operators and geometry of Banach spaces*, NSF-CBMS Regional Conference Series, No. 60, AMS, Providence 1986.

[R] S. Rolewicz, *Metric linear spaces,* Monografie Matematyczne, Tom. 56. [Mathematical Monographs, Vol. 56] PWN-Polish Scientific Publishers, Warsaw, 1972.