

**BENDIXSON'S CRITERION
FOR AUTONOMOUS SYSTEMS WITH
AN INVARIANT LINEAR SUBSPACE**

MICHAEL Y. LI

Dedicated to the memory of Professor G.J. Butler

ABSTRACT. A class of nonlinear autonomous systems ordinary differential equations in \mathbf{R}^n with an invariant linear subspace which includes as examples a wide range of biological and chemical systems is defined and studied. Among other things, criteria precluding the existence of periodic solutions are obtained for such systems using a general method developed in [4].

1. Introduction. Let $D \subset \mathbf{R}^n$ be a convex open set and $x \mapsto f(x) \in \mathbf{R}^n$ a C^1 function defined in D . We consider the autonomous system in \mathbf{R}^n

$$(1.1) \quad x' = f(x)$$

under the following assumptions:

(H1) The Jacobian matrix $\partial f / \partial x$ of the vector field f of (1.1) can be written as

$$(1.2) \quad \frac{\partial f}{\partial x}(x) = -\nu I + A(x) \quad \text{for all } x \text{ in } D,$$

where ν is a constant and $x \mapsto A(x)$ is an $n \times n$ matrix-valued function.

(H2) There exists a constant matrix B with $\text{rank } B = r$ such that

$$(1.3) \quad BA(x) = 0 \quad \text{for all } x \text{ in } D.$$

We will call a nonlinear system (1.1) satisfying (H1) and (H2) an autonomous system with an invariant linear subspace. Examples of

Received by the editors on September 10, 1992, and in revised form on January 19, 1993.

Research is partially supported by the Natural Science and Engineering Research Council of Canada under Grant NSERC A7197 and by the Dissertation Fellowship at the University of Alberta.