Homework 2

(due at 11:00 am on May 3, 2010)

- **Problem 1.** Show that for the gas with Maxwellian distribution the average relative velocity of molecules is of the same order as the average thermal velocity and the speed of sound.
- Problem 2. Derive Bernoulli's equation and its analogue for potential unsteady flow. Discuss the limits of applicability. Use §§5 and 9 of ref. 3.
- **Problem 3.** Under the influence of surface tension σ , a liquid rises to a height H in a glass tube of diameter D. How does H depend on the parameters of the problem? Use dimensional analysis to reveal this dependence.
- **Problem 4.** Construct the solution for inviscid and viscous plane stagnation-point flow, cf. figure 1.



Figure 1: 2D flow near a stagnation point.

Problem 5. Using affine transforms, find the solution for a submerged jet (liquid ejected from a pipe in the space filled with the same liquid, cf. figure 2) in a half-plane $x > 0, -\infty < y < +\infty$:

$$\begin{split} u \, u_x + v \, u_y &= u_{yy}, \\ u_x + v_y &= 0, \\ |u| \to 0, \ y \to \pm \infty. \end{split}$$

Here (u, v) is the velocity field with (x, y)-components, respectively.

Problem 6. Using affine transforms, find the solution for an axisymmetric drop spreading



Figure 2: Submerged jet.

on a flat surface, cf. figure 3, described by the following equation

$$\frac{\partial h}{\partial t} = \frac{2}{3r} \frac{\partial}{\partial r} \left(r h^3 \frac{\partial h}{\partial r} \right),$$

with the boundary condition h = 0 at $r = \infty$ and a mass conservation condition, i.e. mass of the drop should be constant. Make use of a physically relevant conservation law. Determine short and long-time behavior of the solution.



Figure 3: Spreading drop.

Problem 7. Construct self-similar solution for turbulent mixing layer formed by the uniform flow leaving the rectangular body, cf. figure 4.



Figure 4: Turbulent mixing layer.