

Casimir Cascades in Two-Dimensional Turbulence

John C. Bowman (University of Alberta)

Acknowledgements: Jahanshah Davoudi (University of Toronto)

September 8, 2009

`www.math.ualberta.ca/~bowman/talks`

Two-Dimensional Turbulence

- Navier–Stokes equation for **vorticity** $\omega = \hat{\mathbf{z}} \cdot \nabla \times \mathbf{u}$:

$$\frac{\partial \omega}{\partial t} + \mathbf{u} \cdot \nabla \omega = -\nu \nabla^2 \omega + f.$$

- In Fourier space:

$$\frac{\partial \omega_{\mathbf{k}}}{\partial t} = S_{\mathbf{k}} - \nu k^2 \omega_{\mathbf{k}} + f_{\mathbf{k}},$$

where $S_{\mathbf{k}} = \sum_{\mathbf{p}} \frac{\hat{\mathbf{z}} \cdot \mathbf{p} \times \mathbf{k}}{p^2} \omega_{\mathbf{p}}^* \omega_{-\mathbf{k}-\mathbf{p}}^*$.

- When $\nu = 0$ and $f_{\mathbf{k}} = 0$:

energy $E = \frac{1}{2} \sum_{\mathbf{k}} \frac{|\omega_{\mathbf{k}}|^2}{k^2}$ and enstrophy $Z = \frac{1}{2} \sum_{\mathbf{k}} |\omega_{\mathbf{k}}|^2$ are

conserved.

Casimir Invariants

- Inviscid unforced two dimensional turbulence has uncountably many other **Casimir invariants**.
- Any continuously differentiable function of the (scalar) vorticity is conserved by the nonlinearity:

$$\begin{aligned}\frac{d}{dt} \int f(\omega) d\mathbf{x} &= \int f'(\omega) \frac{\partial \omega}{\partial t} d\mathbf{x} = - \int f'(\omega) \mathbf{u} \cdot \nabla \omega d\mathbf{x} \\ &= - \int \mathbf{u} \cdot \nabla f(\omega) d\mathbf{x} = \int f(\omega) \nabla \cdot \mathbf{u} d\mathbf{x} = 0.\end{aligned}$$

- Do these invariants also play a fundamental role in the turbulent dynamics, in addition to the quadratic (energy and enstrophy) invariants? Do they exhibit **cascades**?
- Polyakov [1992] has suggested that the higher-order Casimir invariants cascade to large scales, while Eyink [1996] suggests that they might cascade to small scales.

High-Wavenumber Truncation

- Only the quadratic invariants survive high-wavenumber truncation (Montgomery calls them **rugged invariants**).

$$\frac{\partial \omega_{\mathbf{k}}}{\partial t} = \sum_{\mathbf{p}, \mathbf{q}} \frac{\epsilon_{\mathbf{k}\mathbf{p}\mathbf{q}}}{q^2} \omega_{\mathbf{p}}^* \omega_{\mathbf{q}}^*.$$

where $\epsilon_{\mathbf{k}\mathbf{p}\mathbf{q}} = (\hat{\mathbf{z}} \cdot \mathbf{p} \times \mathbf{q}) \delta(\mathbf{k} + \mathbf{p} + \mathbf{q})$.

- Enstrophy evolution:

$$\frac{d}{dt} \sum_{\mathbf{k}} |\omega_{\mathbf{k}}|^2 = \sum_{\mathbf{k}, \mathbf{p}, \mathbf{q}} \frac{\epsilon_{\mathbf{k}\mathbf{p}\mathbf{q}}}{q^2} \omega_{\mathbf{k}}^* \omega_{\mathbf{p}}^* \omega_{\mathbf{q}}^* = 0.$$

- Invariance of $Z_3 = \int \omega^3 dx$ follows from:

$$0 = \sum_{\mathbf{k}, \mathbf{r}, \mathbf{s}} \left[\sum_{\mathbf{p}, \mathbf{q}} \frac{\epsilon_{\mathbf{k} \mathbf{p} \mathbf{q}}}{q^2} \omega_{\mathbf{p}}^* \omega_{\mathbf{q}}^* \omega_{\mathbf{r}}^* \omega_{\mathbf{s}}^* + 2 \text{ other similar terms} \right].$$

- The absence of an explicit $\omega_{\mathbf{k}}$ in the first term means that setting $\omega_{\ell} = 0$ for $\ell > K$ **breaks the symmetry** in the summations!
- This means that high-wavenumber truncation destroys the invariance of Z_3 .
- However, since the missing terms involve $\omega_{\mathbf{p}}$ and $\omega_{\mathbf{q}}$ for \mathbf{p} and \mathbf{q} higher than the truncation wavenumber K , one might expect almost exact invariance of Z_3 for a well-resolved simulation.
- We will show that this is indeed the case.

Enstrophy Balance

$$\frac{\partial \omega_{\mathbf{k}}}{\partial t} + \nu k^2 \omega_{\mathbf{k}} = S_{\mathbf{k}} + f_{\mathbf{k}},$$

- Multiply by $\omega_{\mathbf{k}}^*$ and integrate over wavenumber angle \Rightarrow enstrophy spectrum $Z(k)$ evolves as:

$$\frac{\partial}{\partial t} Z(k) + 2\nu k^2 Z(k) = 2T(k) + G(k),$$

where $T(k)$ and $G(k)$ are the corresponding angular averages of $\text{Re} \langle S_{\mathbf{k}} \omega_{\mathbf{k}}^* \rangle$ and $\text{Re} \langle f_{\mathbf{k}} \omega_{\mathbf{k}}^* \rangle$.

Nonlinear Enstrophy Transfer Function

$$\frac{\partial}{\partial t} Z(k) + 2\nu k^2 Z(k) = 2T(k) + G(k).$$

- Let

$$\Pi(k) \doteq 2 \int_k^\infty T(p) dp$$

represent the nonlinear transfer of enstrophy into $[k, \infty)$.

- Integrate from k to ∞ :

$$\frac{d}{dt} \int_k^\infty Z(p) dp = \Pi(k) - \epsilon_Z(k),$$

where $\epsilon_Z(k) \doteq 2\nu \int_k^\infty p^2 Z(p) dp - \int_k^\infty G(p) dp$ is the total enstrophy transfer, via dissipation and forcing, **out** of wavenumbers higher than k .

- A positive (negative) value for $\Pi(k)$ represents a flow of enstrophy to wavenumbers higher (lower) than k .
- When $\nu = 0$ and $f_k = 0$:


$$0 = \frac{d}{dt} \int_0^\infty Z(p) dp = 2 \int_0^\infty T(p) dp,$$



so that


$$\Pi(k) = 2 \int_k^\infty T(p) dp = -2 \int_0^k T(p) dp.$$

- Note that $\Pi(0) = \Pi(\infty) = 0$.
- In a steady state, $\Pi(k) = \epsilon_Z(k)$.
- This provides an excellent numerical diagnostic for when a steady state has been reached.


Forcing at $k = 2$, friction for $k < 3$, viscosity for $k \geq k_H = 300$ (1023×1023 dealiased modes)








Cutoff viscosity ($k \geq k_H = 300$)



Cutoff viscosity ($k \geq k_H = 300$)



Molecular viscosity ($k \geq k_H = 0$)

Nonlinear Casimir Transfer

- Fourier decompose the fourth-order Casimir invariant

$$Z_4 = N^3 \sum_j \omega^4(x_j) \text{ in terms of } N \text{ spatial collocation points } x_j:$$

$$Z_4 = \sum_{\mathbf{k}, \mathbf{p}} \omega_{\mathbf{k}} \omega_{\mathbf{p}} \omega_{\mathbf{q}} \omega_{-\mathbf{k}-\mathbf{p}-\mathbf{q}}.$$

$$\begin{aligned} \frac{d}{dt} Z_4 &= \sum_{\mathbf{k}} \left[S_{\mathbf{k}} \sum_{\mathbf{p}, \mathbf{q}} \omega_{\mathbf{p}} \omega_{\mathbf{q}} \omega_{-\mathbf{k}-\mathbf{p}-\mathbf{q}} + 3\omega_{\mathbf{k}} \sum_{\mathbf{p}, \mathbf{q}} S_{\mathbf{p}} \omega_{\mathbf{q}} \omega_{-\mathbf{k}-\mathbf{p}-\mathbf{q}} \right] \\ \frac{d}{dt} Z_4 &= N^2 \sum_{\mathbf{k}} \left[S_{\mathbf{k}} \sum_j \omega^3(x_j) e^{2\pi i \mathbf{j} \cdot \mathbf{k} / N} + 3\omega_{\mathbf{k}} \sum_j S(x_j) \omega^2(x_j) e^{2\pi i \mathbf{j} \cdot \mathbf{k} / N} \right] \\ &\doteq \sum_k T_4(k). \end{aligned}$$


Here $S_{\mathbf{k}}$ is the nonlinear source term in $\frac{\partial}{\partial t} \omega_{\mathbf{k}}$.

Dealiasing: 2/4 Zero Padding Rule

- Computing Z_4 requires computing a double convolution: the Fourier transform of the cubic quantity ω^3 .
- Correctly dealiasing therefore requires a 2/4 zero padding rule (instead of the usual 2/3 rule for a quadratic convolution).


 \Rightarrow even though a 2048×2048 pseudospectral simulation was used, the maximum physical wavenumber retained in each direction was 512.

Downscale Transfer of Z_4



Nonlinear transfer Π_4 of Z_4 averaged over $t \in [250, 740]$.

Vorticity Field




Transfer vs. Flux

- Distinguish between **transfer** and **flux**.
- The mean rate of enstrophy **transfer** to $[k, \infty)$ is given by

$$\Pi(k) = \int_k^\infty T(k) dk = - \int_0^k T(k) dk.$$

- In a steady state, $\Pi(k)$ will trivially be constant within a true inertial range.
- In contrast, the enstrophy **flux** through a wavenumber k is the amount of enstrophy transferred to small scales *via* **triad** interactions involving mode k .

Flux Decomposition for a Single $(\mathbf{k}, \mathbf{p}, \mathbf{q})$ Triad




- Note that energy is conserved: $L_k + S_k = T_k = -T_p - T_q$. Thus

$$L_k = \operatorname{Re} \sum_{\substack{|\mathbf{k}|=k \\ |\mathbf{p}|<k \\ |\mathbf{k}-\mathbf{p}|<k}} M_{\mathbf{k},\mathbf{p}} \omega_{\mathbf{p}} \omega_{\mathbf{k}-\mathbf{p}} \omega_{\mathbf{k}}^* - \operatorname{Re} \sum_{\substack{|\mathbf{k}|=k \\ |\mathbf{p}|<k \\ |\mathbf{k}-\mathbf{p}|>k}} M_{\mathbf{p},\mathbf{k}-\mathbf{p}} \omega_{\mathbf{k}} \omega_{\mathbf{k}-\mathbf{p}} \omega_{\mathbf{p}}^*.$$

Conclusions

- Even though higher-order Casimir invariants do not survive wavenumber truncation, it is possible, with sufficiently well resolved simulations, to check whether they cascade to large or small scales.
- We computed the transfer function of the globally integrated ω^4 inviscid invariant.
- Numerical evidence suggests that in the enstrophy inertial range there is a **direct cascade** of this invariant to small scales.
- However, for the globally integrated ω^3 inviscid invariant, we found no systematic cascade: it appears to slosh back and forth between the large and small scales. This is expected since ω^3 does not have a definite sign.
- One should distinguish between **nonlocal transfer** and **flux**. To compute this decomposition efficiently, one needs to develop a **restricted Fast Fourier transform**.

Asymptote: 2D & 3D Vector Graphics Language



Andy Hammerlindl, John C. Bowman, Tom Prince

<http://asymptote.sf.net>

(freely available under the GNU public license)

Asymptote Lifts T_EX to 3D

$$\int_{-\infty}^{+\infty} e^{-\alpha x^2} dx = \sqrt{\frac{\pi}{\alpha}}$$


<http://asymptote.sf.net>

Acknowledgements: Orest Shardt (U. Alberta)

Kraichnan–Leith–Batchelor Theory

- In an infinite domain
[Kraichnan 1967], [Leith 1968], [Batchelor 1969]:
 - large-scale $k^{-5/3}$ energy cascade;
 - small-scale k^{-3} enstrophy cascade.
- In a **bounded** domain, the situation may be quite different...

Long-Time Behaviour in a Bounded Domain



Tran and Bowman, PRE 69, 036303, 1–7 (2004).

References

- [Batchelor 1969] G. K. Batchelor, Phys. Fluids, **12 II**:233, 1969.
- [Eyink 1996] G. L. Eyink, Physica D., p. 97, 1996.
- [Kraichnan 1967] R. H. Kraichnan, Phys. Fluids, **10**:1417, 1967.
- [Leith 1968] C. E. Leith, Phys. Fluids, **11**:671, 1968.
- [Polyakov 1992] A. Polyakov, 1992, PUPT-1369.
- [Tran & Bowman 2004] C. V. Tran & J. C. Bowman, Physical Review E, **69**:1, 2004.